



## Chemistry Lecture 7: Electrochemistry

Examcrackers MCAT Comprehensive Course, Charles Feng  
[fenguin@gmail.com](mailto:fenguin@gmail.com) / (224) 532-0039

### Redox Reactions

LEO the lion says GER / OIL RIG

Lose electrons = *oxidized*, gain electrons = *reduced*

Something that is reduced is an *oxidizing agent*, if it's oxidized it's a *reducing agent*

Redox reactions involve something being reduced, something being oxidized

### *List of oxidation states:*

Elements=0, F=-1, H=+1 (unless bonded with metal), O=-2 (unless peroxide)

Sum of oxidation states on molecule must equal to charge on molecule

*Balancing:* must balance both MASS and CHARGE.

1. Divide into half reactions (reduction, oxidation)

FOR EACH HALF REACTION...

2. Balance elements that aren't H, O

3. Balance O's by adding H<sub>2</sub>O to one side

4. Balance H's by adding H<sup>+</sup> to one side

5. Balance charge by adding e<sup>-</sup> to one side

NOW COMBINE THEM:

6. Multiply each half reaction by integer so e<sup>-</sup>'s used are equal

7. Add two half reactions together, e<sup>-</sup>'s should cancel out. Also cancel other things if you can.

8. If in basic solution, add OH<sup>-</sup> to both sides until all H's are used up

EXAMPLE  $\text{Cr}(\text{OH})_3(\text{s}) + \text{ClO}_3^-(\text{aq}) \rightarrow \text{CrO}_4^{2-}(\text{aq}) + \text{Cl}^-(\text{aq})$  in basic conditions

### Electric Potential

*Reduction potential:* how much something WANTS to be reduced (generally converting from cation to uncharged element)

*Oxidation potential:* how much something WANTS to be oxidized, equal to negative of reduction potential of oxidized state

All potentials given relative to oxidation potential of H<sub>2</sub> / reduction potential of H<sup>+</sup> = 0

Given something like this:  $\text{Au}^{3+} + 3\text{e}^- \rightarrow \text{Au}(\text{s})$ ,  $E^0 = 1.50 \text{ V}$ :

Reduction potential for Au<sup>3+</sup> is 1.50 V

Oxidation potential for Au (s) is -1.50 V

So Au<sup>3+</sup> is a STRONG oxidizing agent (b/c high reduction potential) but a WEAK reducing agent (b/c low oxidation potential)



## Chemistry Lecture 7: Electrochemistry

Examcrackers MCAT Comprehensive Course, Charles Feng  
[fenguin@gmail.com](mailto:fenguin@gmail.com) / (224) 532-0039

Basically for two compounds reacting together, higher reduction potential one will be reduced, higher oxidation/lower reduction potential one will be oxidized.

Potential of an ionic reaction: add potentials for each *half reaction* (each individual ion)

### Galvanic Cell

AN OX, RED CAT (anode=oxidation, cathode=reduction)

Has an *anode* and a *cathode* connected by a wire where electrons can flow. Oxidation happens at the anode (converting an element to a positive ion) and reduction happens at the cathode (converting an element to a negative ion)

Can figure out which is anode/cathode by looking at reduction potentials: higher reduction potential = will be reduced = cathode

Can have a *salt bridge* where ions in solution can flow to counterbalance the electrons flowing through wire.

*Electromotive force/cell potential*: reduction potential at cathode - oxidation potential at anode (or sum of reduction potentials for elements at cathode + anode)

*Cell diagram*: Anode|AnodeIon||CathodeIon|Cathode

### Formulas

Free energy:  $\Delta G = -nFE$  or  $\Delta G^\circ = -nFE^\circ$ , so positive cell potential = spontaneous reaction

F = Faraday's constant = 96500 C/mol

n = number of moles of electrons in balanced redox reaction

At non-standard state:  $\Delta G = \Delta G^\circ + RT \ln Q$

Can also write  $\Delta G^\circ = -RT \ln K$  where K is equilibrium constant for redox reaction

-> K=1 (equal amount reactants/products):  $\Delta G^\circ = 0$

-> K>1 (more products):  $\Delta G^\circ < 0$  or spontaneous reaction

-> K<1 (less products):  $\Delta G^\circ > 0$  or non spontaneous reaction

### Concentration Cell

Same chemical used for anode/cathode, just at different concentrations

More concentration of positive ion = reduction will occur, or cathode

Less concentration of positive ion = oxidation will occur, or anode

Nernst equation can be used to find cell potential based on concentration difference

### Electrolytic Cell

Supplying power to cause opposite of galvanic cell (backwards from what we expect)